

BRIEFING

Potato Starch. The United States Pharmacopoeia is the coordinating pharmacopoeia for the international harmonization of the compendial standards for the Potato Starch monograph, as part of the process of international harmonization of monographs and general analytical methods of the European, Japanese, and United States pharmacopoeias. The following monograph, which represents the **ADOPTION STAGE 6** document, is based on the corresponding monograph for Potato Starch that was revised based on a request from the Japanese Pharmacopoeia. Differences between the **ADOPTION STAGE 6** document and the current USP monograph for Potato Starch include the following: the sentence under *Identification* test A “Between crossed nicol prisms” is changed to “Between orthogonally oriented polarizing plates or prisms.”

(EM2: K. Moore.)

RTS—C50840

Add the following:

▲ Potato Starch

Attributes	EP	JP	USP
Definition	+	+	+
Identification A	+	+	+
Identification B	+	+	+
Identification C	+	+	+
pH	+	+	+
Loss on Drying	+	+	+
Residue on Ignition	+	+	+
Limit of Iron	+	+	+
Limit of Oxidizing Substances	+	+	+
Limit of Sulfur Dioxide	+	+	+

Legend: + will adopt and implement; – will not stipulate

Nonharmonized attributes: Characters, *Microbiological Examination of Nonsterile Products: Microbial Enumeration Tests and Tests for Specified Microorganisms, Labeling, and Packing and Storage* (USP)

Specific local attributes: Foreign matter (EP)

Reagents and reference materials: Each pharmacopoeia will adapt the text to take account of local reference materials and reagent specifications.

DEFINITION

Potato Starch is obtained from the tuber of *Solanum tuberosum* L.

IDENTIFICATION

- **A. PROCEDURE:** Examine under a microscope using a mixture of equal volumes of glycerol and water.

Acceptance criteria: It presents granules, either irregularly shaped, ovoid or pear-shaped, usually 30–100 µm in size but occasionally exceeding 100 µm, or rounded, 10–35 µm in size. There are occasional compound granules having two to four components. The ovoid and pear-shaped granules have an eccentric hilum and the rounded granules acentric or slightly eccentric hilum. All granules show clearly visible concentric striations. Between orthogonally oriented polarizing plates or prisms, the granules show a distinct black cross intersecting at the hilum.

- **B. PROCEDURE**

Sample solution: 20 mg/mL in water

Analysis: Boil for 1 min, and cool.

Acceptance criteria: A thin, cloudy mucilage is formed.

- **C. PROCEDURE**

Sample solution: 1 mL of the mucilage obtained in *Identification* test B

Analysis: Add 0.05 mL of iodine and potassium iodide TS 2 to the *Sample solution*.

Acceptance criteria: An orange-red to dark blue color is produced, which disappears upon heating.

IMPURITIES

Inorganic Impurities

- **RESIDUE ON IGNITION** 〈 281 〉 : NMT 0.6%, determined on a 1.0-g sample

- **LIMIT OF IRON**

Sample solution: Shake 1.5 g of Potato Starch with 15 mL of 2 N hydrochloric acid, and filter. Transfer 10 mL of the filtrate to a test tube, add 2 mL of citric acid solution (2 in 10), and 0.1 mL of thioglycolic acid. Add 10 N ammonium hydroxide until the solution is distinctly alkaline to litmus, and dilute with water to 20 mL.

Standard iron stock solution A: Equivalent of 10 µg/mL of iron prepared as directed under *Iron* 〈 241 〉

Standard iron stock solution B: 1 µg/mL of iron from *Standard iron stock solution A* in water

[NOTE—Prepare immediately before use.]

Standard iron solution: Transfer 10 mL of *Standard iron stock solution B* to a test tube, and add 2 mL of citric acid solution (2 in 10), and 0.1 mL of thioglycolic acid. Add 10 N ammonium hydroxide until the solution is distinctly alkaline to litmus, and dilute with water to 20 mL.

Acceptance criteria: After 5 min, any pink color in the *Sample solution* is not more intense than that in the *Standard iron solution*, corresponding to a limit of 10 ppm of iron.


● **LIMIT OF SULFUR DIOXIDE**

Carbon dioxide: Use carbon dioxide, with a flow regulator that will maintain a flow of 100 ± 10 mL/min.

Bromophenol blue indicator solution: 0.2 mg/mL of bromophenol blue in dilute alcohol. Filter if necessary.

Hydrogen peroxide solution: Dilute 30% hydrogen peroxide with water to obtain a 3% solution. Just before use, add 3 drops of *Bromophenol blue indicator solution*, and neutralize to a violet-blue endpoint with 0.01 N sodium hydroxide. Do not exceed the endpoint.

Apparatus: *Figure 1.*



In this test, the sulfur dioxide is released from the sample in a boiling acid medium and is removed by a stream of carbon dioxide. The separated gas is collected in a dilute hydrogen peroxide solution where the sulfur dioxide is oxidized to sulfuric acid and titrated with standard alkali. The apparatus consists essentially of a 500-mL three-neck, round-bottom boiling flask, a separatory funnel having a capacity of 100 mL or greater, a gas inlet tube of sufficient length to permit introduction of the carbon dioxide within 2.5 cm of the bottom of the boiling flask, a reflux condenser having a jacket length of 200 mm, and a delivery tube connecting the upper end of the

reflux condenser to the bottom of a receiving test tube. Apply a thin film of stopcock grease to the sealing surfaces of all of the joints except the joint between the separatory funnel and the boiling flask, and clamp the joints to ensure tightness.

Analysis: Add 150 mL of water to the boiling flask. Close the stopcock of the separatory funnel, and begin the flow of carbon dioxide at a rate of 100 ± 5 mL/min through the *Apparatus*. Start the condenser coolant flow. Add 10 mL of *Hydrogen peroxide solution* to a receiving test tube. After 15 min, without interrupting the flow of carbon dioxide, remove the separatory funnel from the boiling flask, and transfer 25.0 g of Potato Starch into the boiling flask with the aid of 100 mL of water. Apply stopcock grease to the outer joint of the separatory funnel, and replace the separatory funnel in the boiling flask. Close the stopcock of the separatory funnel, and add 80 mL of 2 N hydrochloric acid to the separatory funnel. Open the stopcock of the separatory funnel to permit the hydrochloric acid solution to flow into the boiling flask, guarding against the escape of sulfur dioxide into the separatory funnel by closing the stopcock before the last few mL of hydrochloric acid drain out. Boil the mixture for 1 h. Remove the receiving test tube, and transfer its contents to a 200-mL wide-necked, conical flask. Rinse the receiving test tube with a small portion of water, add the rinsing to the 200-mL conical flask, and mix. Heat on a water bath for 15 min, and allow to cool.

Analysis: Add 0.1 mL of *Bromophenol blue indicator solution*, and titrate the contents with 0.1 N sodium hydroxide VS until the color changes from yellow to violet-blue. Perform a blank determination, and make any necessary correction (see *Titrimetry* 〈 541 〉).

Calculate the content, in ppm, of sulfur dioxide in the *Sample solution* taken:

$$\text{Result} = 1000 (32.03) \text{ VN/W}$$

32.03 = milliequivalent weight of sulfur dioxide

V = volume of titrant consumed (mL)

N = normality of the titrant

W = weight of the *Sample solution* (g)

Acceptance criteria: NMT 50 ppm

● LIMIT OF OXIDIZING SUBSTANCES

Sample solution: Transfer 4.0 g to a glass-stoppered, 125-mL conical flask, and add 50.0 mL of water. Insert the stopper, and swirl for 5 min. Transfer to a glass-stoppered, 50-mL centrifuge tube, and centrifuge to clarify. Transfer 30.0 mL of the clear supernatant to a glass-stoppered, 125-mL conical flask. Add 1 mL of glacial acetic acid and 0.5–1.0 g of potassium iodide. Insert the stopper, swirl, and allow to stand for 25–30 min in the dark. Add 1 mL of starch TS.

Analysis: Titrate with 0.002 N sodium thiosulfate VS to the disappearance of the starch–iodine color. Perform a blank determination, and make any necessary correction. Each mL of 0.002 N sodium thiosulfate is equivalent to 34 µg of oxidant, calculated as hydrogen peroxide.

Acceptance criteria: NMT 1.4 mL of 0.002 N sodium thiosulfate is required (20 ppm, calculated as H₂O₂).

SPECIFIC TESTS

- ◆ **MICROBIOLOGICAL EXAMINATION OF NONSTERILE PRODUCTIONS: MICROBIAL ENUMERATION TESTS** 〈 61 〉 and **TESTS FOR SPECIFIED MICROORGANISMS** 〈 62 〉 : The total aerobic microbial count does not exceed 1000 cfu/g; the total combined molds and yeasts count does not exceed 100 cfu/g, and it meets the requirements of the test for the absence of *Escherichia coli*. ◆

- **LOSS ON DRYING** 〈 731 〉 : Dry 1 g, at 130° for 90 min: it loses NMT 20.0% of its weight.

- **pH** 〈 791 〉 : 5.0–8.0

Sample solution: Prepare a slurry by weighing 5.0 g of Potato Starch, transferring to a suitable nonmetallic container, and adding 25.0 mL of freshly boiled and cooled water.

Analysis: Agitate continuously at a moderate rate for 1 min. Stop the agitation, and allow to stand for 15 min. Determine the pH to the nearest 0.1 unit.

ADDITIONAL REQUIREMENTS

- ◆ **PACKAGING AND STORAGE:** Preserve in well-closed containers. No storage requirements specified. ◆

▲ NF28

Auxiliary Information— Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
Monograph	Kevin T. Moore, Ph.D. Senior Scientist 1-301-816-8369	(EM205) Excipient Monographs 2
〈 61 〉	Radhakrishna S Tirumalai, Ph.D. Senior Scientist 1-301-816-8339	(MSA05) Microbiology and Sterility Assurance
〈 62 〉	Radhakrishna S Tirumalai, Ph.D. Senior Scientist 1-301-816-8339	(MSA05) Microbiology and Sterility Assurance

USP32–NF27 Page 1358

Pharmacopeial Forum: Volume No. 35(3) Page 689